Cefiderocol *In Vitro* Activity Against Molecularly Characterized *Acinetobacter baumannii-calcoaceticus* complex and *Pseudomonas aeruginosa* Clinical Isolates Causing Infection in Europe and Adjacent Regions (2020–2021)

R.E. Mendes¹, J.H. Kimbrough ¹, D. Shortridge¹, H.S. Sader¹, M. Castanheira¹ JMI Laboratories, North Liberty, Iowa, USA

Introduction

- Multidrug-resistant (MDR) P a a a and Ac bac ba a -cac ac c complex (A. ba a) cause serious nosocomial infections, especially in intensive care unit patients.
 - These pathogens may be resistant to many clinically available antimicrobial agents, generating therapeutic challenges.
- Cefiderocol is a siderophore-conjugated cephalosporin with broad activity against aerobic, Gram-negative bacteria, including carbapenem-resistant Enterobacterales (CRE), carbapenem-resistant *P. a* a, and *A. ba* a.
 - This cephalosporin utilizes the bacterial iron transport system to gain access to the periplasmic space to reach its targets.
- This study evaluated the activities of cefiderocol and comparator agents against resistant and molecularly characterized
 A. ba a and P. a a recovered from hospitalised patients as part of the SENTRY Antimicrobial Surveillance Program for Europe and surrounding regions.

Results