IDWeek 2022 | Poster #P642

I V Activity of Cefiderocol and Comparator Agents Against Molecularly Characterized Clinical Isolates of Enterobacterales Causing Infections in United States Hospitals (2020–2021)

RE Mendes, JH Kimbrough, V Kantro, D Shortridge, HS Sader, JM Streit, M Castanheira JMI Laboratories, North Liberty, Iowa, USA

Introduction

- Cefiderocol is approved by the US Food and Drug Administration (FDA)
 for the treatment of complicated urinary tract infections, including
 pyelonephritis, as well as hospital-acquired bacterial pneumonia, and
 ventilator-associated bacterial pneumonia.
- Cefiderocol is a siderophore cephalosporin with broad activity against Gram-negative bacteria, including multidrug-resistant (MDR) organisms like carbapenem-resistant Enterobacterales (CRE), carbapenem-resistant P e, domona ae , gino a, and Acine obac e ba, mannii.
- The activity of this molecule is due to its ability to achieve high periplasmic concentrations by hijacking the bacterial iron transport machinery, which in turn potentiates cell entry.
- In addition, cefiderocol remains stable to hydrolysis by serine
 -lactamases (ESBLs, KPCs, and OXA-type carbapenemases) and metallo- -lactamases.
- In this study, the activities of cefiderocol and comparator agents were analyzed against Enterobacterales, including molecularly characterized isolates, as part of the SENTRY Antimicrobial Surveillance Program for USA.

Materials and Methods

Bacterial organisms

- This study comprised a collection of 8,328 Enterobacterales collected from various clinical specimens from patients hospitalized in 32 medical centers in all 9 US Census Divisions during 2020–2021. Only consecutive isolates (1 per patient infection episode) responsible for documented infections according to local criteria were included.
- Bacterial identification was confirmed by standard algorithms supported by matrix-assisted laser desorption ionization-time of flight mass spectrometry (Bruker Daltonics, Bremen, Germany).

Susceptibility testing

- Isolates were tested for susceptibility by broth microdilution following the Clinical and Laboratory Standards Institute (CLSI) M07 (2018) guidelines.
- Frozen-form broth microdilution panels were manufactured by JMI
 Laboratories (North Liberty, IA, USA) and contained cation-adjusted MuellerHinton broth for comparator agents.
- Susceptibility testing for cefiderocol used broth microdilution panels containing iron-depleted media per CLSI guidelines.
- Quality assurance was performed by sterility checks, colony counts, and testing CLSI-recommended quality control reference strains.
- MIC interpretations were performed using CLSI breakpoints for comparators and FDA/CLSI breakpoints for cefiderocol (4/8/16 mg/L for susceptible, intermediate, and resistant).
- Imipenem-relebactam MIC interpretations used FDA breakpoints.

 E che ichia coli, Kleb iella ne moniae, and P o e mi abili with ceftriaxone, ceftazidime, or aztreonam MIC values of 2 mg/L and any Enterobacterales displaying MIC values 2 mg/L for imipenem (excluded for P. mi abili, P. enne i, and indole-positive Proteeae) or meropenem were subjected to genome sequencing and screening of -lactamase genes.

Screening of -lactamase genes

- Selected isolates had total genomic DNA extracted by the fully automated
 Thermo Scientific™ KingFisher™ Flex Magnetic Particle Processor (Cleveland,
 OH, USA), which was used as input material for library construction.
- DNA libraries were prepared using the Nextera™ library construction protocol (Illumina, San Diego, CA, USA) following the manufacturer's instructions and were sequenced on MiSeq Sequencer platforms at JMI Laboratories.
- FASTQ format sequencing files for each sample set were assembled independently using *de no o* assembler SPAdes 3.15.3. An in-house software was applied to align the assembled sequences against a comprehensive in-house database containing known -lactamase genes.

Conclusions

- Cefiderocol in i o activity (98.4% susceptible) was consistent against various subsets, including against Enterobacterales carrying carbapenemase genes other than bla_{KPC}, against which approved -lactam/ -lactamase inhibitor combinations showed limited activity.
- These data reinforce cefiderocol as an important option for the treatment of serious infections caused by Enterobacterales and resistant subsets in patients hospitalized in US medical centers.

Acknowledgments

This research and poster presentation were sponsored by Shionogi & Co., LTD.

References

- 1. Clinical and Laboratory Standards Institute. 2018. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. *M07* 11 h Edi ion. Wayne, PA, USA.
- 2. Clinical and Laboratory Standards Institute. 2022. Performance standards for antimicrobial susceptibility testing. *M100 32nd Edi ion*. Wayne, PA, USA.
- 3. FDA Susceptibility Test Interpretive Criteria: https://www.fda.gov/drugs/development-resources/antibacterial-susceptibility-test-interpretive-criteria. Accessed April 2022.
- 4. Mendes RE, Jones RN, Woosley LN, Cattoir V, Castanheira M. 2019. Application of next-generation sequencing for characterization of surveillance and clinical trial isolates: Analysis of the distribution of -lactamase resistance genes and lineage background in the United States. O en Fo : m Irf ec Di 6: S69-S78.
- 5. Ong'uti S, Czech M, Robilotti E, Holubar M. 2021. Cefiderocol: A new cephalosporin stratagem against multidrug resistant Gram-negative bacteria. Clin Irf ec Di . In e .
- 6. Syed YY. 2021. Cefiderocol: A review in serious Gram-negative bacterial infections. D g . 24: 1–13.

Contact

Rodrigo E. Mendes, Ph.D.

JMI Laboratories


345 Beaver Kreek Centre, Suite A

North Liberty, Iowa 52317

Phone: (319) 665-3370

Fax: (319) 665-3371

Email: rodrigo-mendes@jmilabs.com



is stored.

visit https://www.jmi labs.com/data/posters /IDWeek2022_Ce der ocolVsMolecularEnteros .pdf

No personal information